

T=1 and T=0 states in the N=Z=43 nucleus, $^{86}\text{Tc}^*$

A. B. Garnsworthy^{1,2}, P. H. Regan¹, S. Pietri¹, D. Rudolph³, Zs. Podolyák¹, S. J. Steer¹, F. Becker⁴, P. Bednarczyk⁴, J. Gerl⁴, M. Górska⁴, H. Grawe⁴, I. Kojouharov⁴, H. Schaffner⁴, H. J. Wollersheim⁴, J. Grębosz^{5,4}, G. Benzoni⁶, B. Blank⁷, C. Brandau¹, A. M. Bruce⁸, L. Caceres^{4,9}, F. Camera⁶, W. N. Catford¹, I. J. Cullen¹, Zs. Dombrády¹⁰, P. Doornenbal⁴, E. Estevez¹¹, H. Geissel⁴, W. Gelletly¹, A. Heinz², R. Hoischen³, G. Ilie¹², J. Jolie¹², G. A. Jones¹, A. Jungclaus⁹, A. Kelic⁴, M. Kmiecik⁵, F. G. Kondev¹³, T. Kurtukian-Nieto¹¹, N. Kurz⁴, S. Lalkovski¹⁴, Z. Liu¹, A. Maj⁵, S. Myalski⁵, F. Montes⁴, M. Pfützner¹⁵, T. Saito⁴, T. Shizuma^{1,16}, A. J. Simons¹, S. Schwertel¹⁷, S. Tachenov⁴, P. M. Walker¹, E. Werner-Malento^{4,15}, O. Wieland⁶

¹Department of Physics, University of Surrey, Guildford, GU2 7XH, UK

²WNSL, Yale University, 272 Whitney Avenue, New Haven, CT, 06520, USA

³Department of Physics, Lund University, S-22100, Lund, Sweden

⁴GSI, Planckstrasse 1, D-64291, Darmstadt, Germany

⁵The Henryk Niewodniczański Institute of Nuclear Physics, PL-31-342, Kraków, Poland

⁶Università degli Studi di Milano and INFN sezione di Milano, I-20133, Milano, Cmerino, Italy

⁷CENBG, le Haut Vigneau, F-33175, Gradignan Cedex, France

⁸School of Engineering, University of Brighton, Brighton, BN2 4GJ, UK

⁹Departemento di Teórica, Universidad Autonoma de Madrid, E-28049, Madrid, Spain

¹⁰Institute for Nuclear Research, H-4001, Debrecen, Hungary

¹¹Universidad de Santiago de Compostela, E-15706, Santiago de Compostela, Spain

¹²IKP, Universität zu Köln, D-50937, Köln, Germany

¹³Nuclear Engineering Division, Argonne National Laboratory, Argonne, IL-60439, USA

¹⁴Faculty of Physics, University of Sofia "St. Kliment Ohridsk" Sofia, Bulgaria

¹⁵IEP, Warsaw University, Hoża 69, PL-00-681, Poland

¹⁶Japan Atomic Energy Research Institute, Kyoto, 619-0215, Japan and

¹⁷Physik Department E12, Technische Universität München, Garching, Germany

The low-lying structure of ^{86}Tc has been studied using isomer-decay spectroscopy at GSI in the first experiment of the Stopped-Beam phase within the Rare ISotope INvestigation at GSI (RISING) campaign. Following projectile fragmentation of a 750 MeV/u beam of ^{107}Ag , reaction products were separated and unambiguously identified using the FRagment Separator (FRS) in combination with its ancillary detectors. The ions were made to stop in a plastic stopper at the final focus of the FRS in the centre of the Stopped RISING γ -ray spectrometer. This high-efficiency, high-granularity array consists of 15 germanium cluster detectors in a compact configuration which provides a full photopeak efficiency in excess of 15% at 1.3 MeV. Internal decay of the previously identified [1] microsecond isomer in ^{86}Tc was confirmed with the addition of two previously unobserved γ -rays which help to determine the excitation energy and spin of the isomeric state. As in other heavy odd-odd N=Z nuclei a notably lower density of states below 1 MeV excitation energy compared with neighbouring odd-odd nuclei away from the N=Z line [2] is observed with the lowest T=0 state identified in the preliminary analysis being a $J^\pi=3^+$ state located 1176 keV above the T=1 [3] groundstate. Results from this experiment will be discussed along with assignments of structure made from shell model calculations and systematics of N=Z nuclei.

[1] C. C. Chandler *et al.*, Phys. Rev. C 61, 044309 (2000)

[2] D. J. Jenkins, *et al.*, Phys. Rev. C 65, 064307 (2002)

[3] C. Longour, *et al.*, Phys. Rev. Lett. 81, 3337 (1998)

* This work was partially supported by the EPSRC (UK) and the U. S. Department of Energy under grants DE-FG02-91ER-40609 and W-31-109-ENG-38. ABG would also like to acknowledge financial support from Nexia Solutions Limited, a subsidiary of BNFL.